

FDLquest #3: Dermatology

Compiled by Brayden Zaharia

Edited by Ella Shi, Yashmeet Kaur, Trevor Chan, Sophia
Sung and Gloria Ng

35 slides

ALL AGENTS CONTRAINDICATED IN HYPERSENSITIVITY UNLESS OTHERWISE STATED

Benzoyl Peroxide

Acne Vulgaris Therapy

Brand name(s):



Doses/forms:

Benzagel:

Gel:



2.5%



5%

Wash:



5%

Clean & Clear Persa-Gel:

Lotion:



5%

Indication(s):

Indicated for the topical treatment of acne vulgaris

Benzac:

Gel:



10%

PanOxyl:

Cream:



4%

Tretinoin

Acne Vulgaris Therapy

Brand name
D/C

Brand name(s):

Stieva-A[®]

Retin-A[®]
(tretinoin)

RETIN-A MICRO[®]
(tretinoin) Gel microsphere

Vitamin A acid

Doses/forms:

Stieva-A
Cream:



0.01%



0.025%



0.05%

Retin-A Micro
Gel:



0.04%



0.1%

Indication(s):

Indicated for the topical treatment
of acne vulgaris

Contraindication(s):

Tretinoin is contraindicated in
pregnancy and lactation

Retin-A:

Cream:



0.05%

Gel:



0.025%



0.01%

Vitamin A Acid
Gel:



0.025%



0.05%

Adapalene

Acne Vulgaris Therapy

Brand name(s):

DIFFERIN® 

Indication(s):

Indicated for the topical treatment of acne vulgaris

Contraindication(s):

Adapalene is contraindicated in those with eczema or seborrheic dermatitis

Doses/forms:

Differin:

Cream:



0.1%

Gel:



0.1%

Differin XP

Gel:



0.3%

Tazarotene

Antipsoriatic, Acne Vulgaris Therapy

Brand name(s):

TAZORAC®

Discontinued

ARAZLO®
(tazarotene) Lotion, 0.045%

Indication(s):

Indicated for the topical
treatment of acne vulgaris

Tazarotene is also indicated for
treatment of psoriasis and
premature aging

Doses/forms:

Tazorac

Gel:



0.1%

Cream:



0.05%

Arazlo

Lotion:



0.045%

Clindamycin phosphate / benzoyl peroxide

Acne Vulgaris Therapy

Brand name(s):



Doses/forms:

BenzaClin, Clindoxyl
Gel:



1% Clindamycin
5% Benzoyl Peroxide

Clindoxyl ADV
Gel:



1% Clindamycin
3% Benzoyl Peroxide

Indication(s):

Indicated for the topical treatment
of acne vulgaris

Adapalene / benzoyl peroxide

Acne Vulgaris Therapy

Brand name(s):



Indication(s):

Indicated for the topical treatment of acne vulgaris

Contraindication(s):

Adapalene is contraindicated in those with eczema or seborrheic dermatitis

Doses/forms:

TactuPump
Gel:



0.1% Adapalene
+2.5% Benzoyl
Peroxide

TactuPump Forte
Gel:



0.3% Adapalene
+2.5% Benzoyl
Peroxide

Clindamycin / tretinoin

Acne Vulgaris Therapy

Brand name(s):

biacna

Doses/forms:

Gel:



Indication(s):

Indicated for the topical treatment of acne vulgaris

Contraindication(s):

Tretinoin is contraindicated in pregnancy and lactation

0.025% Tretinoin
+1.2% Clindamycin

Erythromycin / benzoyl peroxide

Acne Vulgaris Therapy

Brand name(s):

Benzamycin®

Doses/forms:

Gel:



Indication(s):

Indicated for the topical treatment of acne vulgaris

3% Erythromycin
+5% Benzoyl Peroxide

Isotretinoin

Acne Vulgaris Therapy

Brand name(s):



epuris®



Indication(s):

Isotretinoin is an oral agent used in the treatment of severe nodular/inflammatory acne vulgaris

Contraindication(s):

Isotretinoin is contraindicated in:

1. Pregnancy and lactation
2. Hepatic and renal insufficiency
3. Elevated blood lipid values
4. Patients taking tetracyclines

Doses/forms:

Accutane, Clarus:

Caps:

10mg



40 mg



Epuris:

Caps:

10mg



20mg



30mg



40mg



Absorica LD:

Caps:

8mg



16mg



24mg



32mg



Acitretin

Topical Calcineurin Inhibitor

Doses/forms:


Brand name(s):

^{Pr} **SORIATANE**[®]

Indication(s):

Indicated for the topical treatment of
acne vulgaris

Caps:



10 mg



25 mg

Tacrolimus

Topical Calcineurin Inhibitor

Brand name(s):



Indication(s):

Tacrolimus and Pimecrolimus are indicated for treatment and maintenance of atopic dermatitis

Doses/forms:

Ointment:



0.03% w/w



0.1% w/w

Pimecrolimus

Topical Calcineurin Inhibitor

Brand name(s):



Indication(s):

Tacrolimus and Pimecrolimus are indicated for treatment and maintenance of atopic dermatitis

Doses/forms:

Cream:



1%

Ivermectin

Topical Rosacea Therapy

Brand name(s):



Doses/forms:

Cream:



Indication(s):

Ivermectin is indicated for treatment of inflammatory lesions of rosacea in adults (≥ 18 years old)

1%

Brimonidine (Tartrate)

Topical Rosacea Therapy

Brand name(s):



Indication(s):

Brimonidine is indicated for treatment of facial erythema of rosacea in adults (>18 years old)

Doses/forms:

Gel:



0.33%

Azelaic acid

Topical Rosacea Therapy

Brand name(s):



Doses/forms:

Gel:



15% w/w

Indication(s):

Indicated for the topical treatment of
acne vulgaris

Spironolactone

Aldosterone receptor antagonist

Brand name(s):

Aldactone®
spironolactone

Indication(s):

Indicated for the topical treatment of
acne vulgaris

Doses/forms:

Tab:



25 mg



100 mg

Corticosteroids

The diagram below lists the number of different types of dosage forms available for each class of steroid & route. For example, mometasone furoate has 3 types of topical dosage forms available: cream, lotion and ointment.

Drug	Nasal	Ophth	Syst.	Topical
Fluticasone propionate	✓			
Mometasone furoate monohydrate	✓			
Prednisolone acetate		✓		
Prednisone			✓	
Dexamethasone		2	3	
Hydrocortisone			✓	3 (+2 rectal forms)
Betamethasone Valerate				4
Betamethasone dipropionate				3
Clobetasol 17-propionate				5
Triamcinolone acetonide	✓		✓	3
Mometasone furoate				3

INDICATIONS BY ROUTE:

NASAL: all are Indicated for seasonal and perennial allergies

OPHTHALMIC: Indicated for ocular inflammation

SYSTEMIC: Indicated for endocrine disorders as well as severe inflammatory conditions such as allergic conditions and asthma

TOPICAL: Indicated for inflammatory skin conditions

Fluticasone propionate

Nasal corticosteroid

Brand name(s):



Indication(s):

Indicated for seasonal and perennial allergies

Doses/forms:

Suspension:



50 µg/metered spray

Mometasone furoate monohydrate

Nasal/Topical corticosteroid

Brand name(s):

Doses/forms:

Nasonex[®]
(Mometasone Furoate Monohydrate)

Suspension (Nasonex):

(Elocom):

Elocom[®]
mometasone furoate

Cream:

Oint:

Lotion:



Indication(s):

50 µg/metered spray

ALL 0.1%

- Nasonex indicated for seasonal and perennial allergies
- Elocom indicated for inflammatory skin conditions

Prednisolone acetate

Ophthalmic Corticosteroid

Brand name(s):



Doses/forms:

Suspension:



1% w/v

Indication(s):

Indicated for ocular
inflammation

Contraindication(s):

Ophthalmic corticosteroids are
contraindicated in ocular herpes
simplex and advanced glaucoma.

Prednisone

Systemic corticosteroid

Brand name(s):

Prednisone

(brand name d/c)



Indication(s):

Indicated for endocrine disorders as well as severe inflammatory conditions such as allergic conditions and asthma

Doses/forms:

Prednisone

Tabs:



5 mg



50 mg

Winpred

Tabs:



1 mg

Dexamethasone (systemic)

Systemic corticosteroid

Brand name(s):

- See Dexamethasone (ophthalmic)

Indication(s):

Indicated for endocrine disorders as well as severe inflammatory conditions such as allergic conditions and asthma

Doses/forms:

Tabs:



0.5mg



0.75mg



2mg



4mg

Elixir:



0.5 mg/5 mL

Injection:



4mg/mL




10mg/mL

Dexamethasone (Ophthalmic)

Ophthalmic corticosteroid

Brand name(s):



Doses/forms:

Ophthalmic oint: Ophthalmic susp:



0.1% w/w



0.1% w/v

Indication(s):

Indicated for ocular inflammation

Contraindication(s):

Ophthalmic corticosteroids are contraindicated in ocular herpes simplex and advanced glaucoma.

Hydrocortisone

Topical/Systemic corticosteroid

Brand name(s):



HydermTM



CORTATE[®]



SOLU-CORTEF[®]



Cortoderm[®]
Hydrocortisone Ointment USP, 0.5%
Onguent d'hydrocortisone USP 0,5% 15 g



CORTEF
10 mg HYDROCORTISONE

Indication(s):

Systemic: endocrine disorders as well as severe inflammatory conditions such as allergic conditions and asthma

Topical: inflammatory skin conditions (available for MAC)

Hydrocortisone (Cont.)

Topical/Systemic corticosteroid


Doses/forms:

<u>Cream:</u>		<u>Pwd for inj:</u>	<u>Lotion:</u>
<u>Cortate</u>	<u>Generic</u>	<u>(Solu-Cortef)</u>	<u>(Generic)</u>
	  	 	
0.5%	0.2% 1% 2.5%	0.1g 0.25g	1%
	<u>Hyderm</u>	 	
	 	0.5g 1g	
	0.5% 1%		
<u>Ointment:</u>		<u>Tablets:</u>	
<u>(Cortoderm)</u>		<u>(Cortef)</u>	
 		 	
0.5% 1%		10mg 20mg	

Betamethasone Valerate

Topical Corticosteroid

Brand name(s):



Discontinued

Indication(s):

Indicated for inflammatory skin conditions

Doses/forms:

Betaderm:

Cream:



0.05%



0.1%

Scalp lotion:



0.1%

Ointment:



0.05%



0.1%

Clobetasol 17-propionate

Topical Corticosteroid

Brand name(s):

Clobex
(clobetasol propionate) Spray, 0.05%

Dermovate
clobetasol propionate

Indication(s):

Indicated for inflammatory skin conditions

Doses/forms: **ALL 0.05%**

Dermovate:

Cream:



Ointment:



Scalp Lotion:



Clobex:

Shampoo: Spray:





Triamcinolone acetonide

Nasal, Topical & Systemic Corticosteroid

Brand name(s):



Discontinued



Doses/forms:

Nasacort Suspension:



55 µg/metered spray

Oracort Dental Paste:



0.1%

Kenalog Injectable Susp:



10mg/mL



40 mg/mL

Indication(s):

Nasal: Indicated for seasonal and perennial allergies

Systemic: endocrine disorders as well as severe inflammatory conditions such as allergic conditions and asthma

Topical: Indicated for inflammatory skin conditions

Aristocort-R Cream: Ointment:



0.1%



0.1%

Aristocort-C Cream:



0.5%

Betamethasone dipropionate


Topical Corticosteroid

Brand name(s):



Doses/forms: ALL 0.05%

Cream:



Ointment:



Lotion:



Indication(s):

Indicated for inflammatory skin conditions

Calcipotriol

Topical Nonsteroidal Antipsoriatic – Vitamin D Analogue

Brand name(s):



Doses/forms:

Ointment:



50 µg/g Calcipotriol

Indication(s):

Used as Topical Nonsteroidal
Anti psoriatic vitamin D
analogue

Calcipotriol / Betamethasone dipropionate

Topical Nonsteroidal Antipsoriatic - Vitamin D Analogue - Corticosteroid

Brand name(s):

Dovobet®

Enstilar®

Doses/forms:

Dovobet

Gel:



Ointment:



Enstilar

Foam:



Generic

Gel:




ALL 50 µg/g Calcipotriol +
0.5mg/g Betamethasone

Permethrin

Lice Treatment - Scabies Treatment

Brand name(s):



Indication(s):

1% permethrin is indicated in the treatment of **lice** infestation

5% permethrin is indicated in the treatment of **scabies** infestation

Contraindication(s):

Permethrin is contraindicated in hypersensitivity to chrysanthemum and ragweed

Doses/forms:

For Lice:

Nix Creme Rinse

Creme Rinse:



1% w/w

For Scabies:

Nix Dermal Cream

Dermal Cream:



5% w/w

Kwellada-P Creme Rinse

Crème Rinse:



1% w/w

Kwellada-P Lotion

Lotion:



5% w/v

Pyrethrins / Piperonyl butoxide

Lice Treatment

Brand name(s):



Indication(s):


Indicated in the treatment of lice infestation

Contraindication(s):

Pyrethrins are contraindicated in hypersensitivity to chrysanthemum, ragweed **and** petrolatum

Doses/forms:

Shampoo:



0.33% w/v Pyrethrins
3% w/v Piperonyl Butoxide

Isopropyl myristate/cyclomethicone

Lice Treatment

Brand name(s):

Resultz

Indication(s):

Indicated in the treatment of lice infestation

Doses/forms:

Topical Solution:



50% w/w Isopropyl Myristate
+50% w/w Cyclomethicone

Dimeticone/Dimethicone

Lice Treatment

Brand name(s):



Indication(s):

Indicated in the treatment of lice infestation

Doses/forms:

Topical solution:



50% w/w



**Help us make the next one even
better!**

- Email ptalc.ubc@gmail.com to let us know how we can improve these slides for your learning.